

Angiología



Nota Técnica

A hybrid clampless technique for aortoiliac derivation as a preparation for renal transplantation

Técnica híbrida clampless para derivación aortoilíaca como preparación para trasplante renal

Xavier Faner Capó, Marvin E. García Reyes, Anna Gené Mola, Lluís Sánchez Besalduch, David Flota Ruiz, Sergi Bellmunt Montoya *Angiology, Vascular Surgery and Endovascular Department. Hospital Universitari Vall d'Hebron. Barcelona, Spain*

Abstract

Introduction: renal transplantation is the treatment of choice for end-stage renal failure. Different aortoiliac reconstruction techniques have proven feasible in preparing for renal transplantation in patients with severe aortoiliac atherosclerosis. However, anatomies with heavy circumferential calcifications or shaggy arteries with extensive thrombus can themselves be a technical contraindication for these reconstructions. The aim of the work is to present a hybrid technique of ilio-femoral bypass performing a proximal anastomosis without clamping the donor artery, serving as an option for renal transplant preparation in these patients.

Material and methods: we present the case of a 73-year-old male with end-stage renal failure deemed unsuitable for renal transplantation due to severe aortoiliac atherosclerosis. After preoperative computed tomography evaluation, he was considered eligible for a left ilio-femoral bypass using a clampless hybrid technique. Following dissection of the left femoral tripod, the left common iliac artery was exposed. A small non-calcified window suitable for direct puncture was found in the artery, through which a guidewire was advanced into the aorta under fluoroscopic guidance. Then, a long over-the-wire sheath with the bevelled prosthetic graft around it was inserted. The graft was sutured to the adventitial layer without any arteriotomy or clamping, mimicking the final aspect of a side-to-end anastomosis. The anastomosis was dilated with a balloon and protected with a covered stent from de graft to the aorta. After tunnelling the graft to the femoral artery, the distal anastomosis was performed in the usual fashion.

Results: this technique was successfully used, and the procedure remained patent 10 months later, without any complications related to the surgery. The patient is still waiting for the transplantation.

Conclusions: a hybrid anastomosis technique without clamping is a feasible option in patients with prohibitive atherosclerosis. Long-term durability and transplant outcomes are still unknown.

Keywords: Hybrid clampless technique. Aortoiliac derivation. Renal transplantation.



http://www.capitulodeflebologia.org/pdf/sobre_tus_venas.pdf

Received: 08/01/2023 • Accepted: 08/01/2023

Conflicts of interest: the authors declare no conflict of interest.

Artificial intelligence: the authors declare not to have used artificial intelligence (Al) or any Al-assisted technologies in the elaboration of the article.

Faner Capó X, García Reyes ME, Gené Mola A, Sánchez Besalduch L, Flota Ruiz D, Bellmunt Montoya S. A hybrid clampless technique for aortoiliac derivation as a preparation for renal transplantation. Angiología 2024;76(5):343

DOI: http://dx.doi.org/10.20960/angiologia.00617

Correspondence:

Marvin E. García Reyes. Angiology, Vascular Surgery and Endovascular Department. Hospital Universitari Vall d'Hebron. Passeig Vall d'Hebron, 119-129. 08035 Barcelona, Spain

e-mail: marvinernesto.garcia@vallhebron.cat